

Galton–Watson branching processes and the growth of gravitational clustering

Ravi K. Sheth

Berkeley Astronomy Department, University of California, Berkeley, CA 94720

1 February 2008

ABSTRACT

The Press–Schechter description of gravitational clustering from an initially Poisson distribution is shown to be equivalent to the well studied Galton–Watson branching process. This correspondence is used to provide a detailed description of the evolution of hierarchical clustering, including a complete description of the merger history tree. The relation to branching process epidemic models means that the Press–Schechter description can also be understood using the formalism developed in the study of queues. The queueing theory formalism, also, is used to provide a complete description of the merger history of any given Press–Schechter clump. In particular, an analytic expression for the merger history of any given Poisson Press–Schechter clump is obtained. This expression allows one to calculate the partition function of merger history trees. It obeys an interesting scaling relation; the partition function for a given pair of initial and final epochs is the same as that for certain other pairs of initial and final epochs.

The distribution function of counts in randomly placed cells, as a function of time, is also obtained using the branching process and queueing theory descriptions. Thus, the Press–Schechter description of the gravitational evolution of clustering from an initially Poisson distribution is now complete. All these interrelations show why the Press–Schechter approach works well in a statistical sense, but cannot provide a detailed description of the dynamics of the clustering particles themselves. One way to extend these results to more general Gaussian initial conditions is discussed.

Key words: galaxies: clustering – galaxies: evolution – galaxies: formation – cosmology: theory – dark matter.

1 INTRODUCTION

This paper is mainly concerned with providing a complete description of gravitational clustering from an initially Poisson distribution. This is not because one thinks it likely that the initial conditions for gravitational clustering in our universe were Poisson. Indeed, measurements of the spectrum of temperature fluctuations in the microwave background suggest otherwise. At present, however, analytic understanding of the growth of clustering from more general Gaussian initial conditions is not as detailed as for the Poisson case studied here. Thus, the Poisson model serves as a convenient toy model with which to study the evolution of nonlinear clustering. Many of the results obtained in this paper should provide at least qualitative insight into the evolution of clustering from more general initial conditions.

The Press–Schechter approach allows one to estimate the distribution of the masses of virialized clumps at a given epoch directly from the initial density distribution. It is based on the hypothesis that, provided the initial velocities are sufficiently small (i.e., that the initial field is sufficiently

cold), overdense regions in the initial density field will eventually collapse to form nonlinear structures. Thus, by studying the statistics of overdense regions in the initial field, one can infer properties of the nonlinear distribution (Press & Schechter 1974; Bond *et al.* 1991; Lacey & Cole 1993). One of the main simplifying assumptions of the Press–Schechter approach is that the overdense regions are assumed to collapse spherically. While this may be a good approximation in the mean, N -body simulations of gravitational clustering from initially scale-free Gaussian random fields show that the collapse of overdense regions is seldom exactly spherical. Nevertheless, the Press–Schechter mass functions for these initially scale-free Gaussian fields have been shown to be in good agreement with the distribution of clump sizes that are measured in relevant N -body simulations (Efstathiou *et al.* 1988; Lacey & Cole 1994).

The Press–Schechter excursion set description of clustering from an initially Poisson distribution of identical particles has been derived (Epstein 1983; Sheth 1995). The probability that a Poisson Press–Schechter clump has N particles is

$$\eta(N, b) = \frac{(Nb)^{N-1} e^{-Nb}}{N!}, \quad (1)$$

where $N \geq 1$ and $0 \leq b < 1$, and b is related to the Press–Schechter overdensity threshold δ_c :

$$b = 1/(1 + \delta_c). \quad (2)$$

For an initially cold Poisson distribution, the threshold overdensity δ_c decreases as the Universe expands in such a way that $b = 0$ initially, and $b \rightarrow 1$ as the clustering develops. If clumps collapse spherically in a universe with critical density, and both growing and decaying modes are present in the linear perturbation theory, then $\delta_c = (5/3) 1.69/a$, where a is the expansion factor (e.g. Bond *et al.* 1991; Lacey & Cole 1993). Thus, b changes rapidly at first, but at late times its evolution is slowed by the expansion of the Universe. (See section 2.3 in Sheth 1995 for another description of the evolution of b that shows this same behaviour.) Since b is known to increase monotonically with time, it will be treated as a pseudo-time variable in the remainder of this paper.

Equation (1) is known as a Borel distribution (Borel 1942). Epstein (1983) derived this distribution by studying the properties of level excursions of a Poisson distribution. His approach is the discrete analog of that which was used later by Bond *et al.* (1991) in their analysis of the initially Gaussian random fields. The Borel distribution can also be derived using simple ‘cloud-in-cloud’ conditional probabilities for a Poisson distribution (Sheth 1995). This approach is the discrete analog of Jedamzik’s (1995) treatment of the Gaussian case.

If the probability that a randomly chosen clump contains N particles is a Borel distribution, then the probability that a randomly chosen particle is in such a clump is $N\eta(N, b)/\langle N \rangle = (1 - b)N\eta(N, b)$, since the average number of particles in a Borel clump is $\langle N \rangle = 1/(1 - b)$. In the limit of large N and small δ_c , Stirling’s approximation for the factorial term implies that

$$\begin{aligned} (1 - b) N \eta(N, b) &= \frac{\delta_c}{(1 + \delta_c)} \left(\frac{N}{1 + \delta_c} \right)^{N-1} \frac{e^{-N/(1+\delta_c)}}{(N-1)!} \\ &\rightarrow \frac{\delta_c}{\sqrt{2\pi N}} \exp\left(-\frac{N\delta_c^2}{2}\right) \end{aligned} \quad (3)$$

(Epstein 1983; Sheth 1995). The final expression is precisely that which obtains for a Gaussian density field with white noise initial fluctuations (e.g. Bond *et al.* 1991). This shows that the Poisson distribution studied in the remainder of this paper can be thought of as the discrete analog of the white noise Gaussian studied by Bond *et al.* (1991), and by Lacey & Cole (1993, 1994).

Sections 2 and 3 study other derivations of the Borel distribution. These derivations provide new insight into the accuracy and applicability of the Press–Schechter approach. An analytic expression that is, essentially, the partition function that describes all possible merger histories of a given Press–Schechter clump is derived in Section 3. Its properties are consistent with those inferred previously (Sheth 1995). In particular, it is consistent with the physical requirement that, in the limit of very small time steps, the probability that a clump has two progenitors should be an infinitesimal of smaller order than the probability that it has three, or more, progenitors. Section 3 also shows that the partition function, and so the growth of clustering, satisfies an

interesting scaling relation. In Section 4, this scaling relation is exploited to provide insight into the details of the growth of hierarchical clustering. Section 4 also shows that this Poisson Galton–Watson, Press–Schechter description is in qualitative agreement with the results of the numerical algorithm developed by Kauffmann & White (1993). Section 5 discusses ways in which the branching process extension of the Press–Schechter approach that is developed in this paper can be extended to provide a detailed description of clustering from initially Gaussian random fields. Appendices A, B and C provide details of some of the calculations.

Appendix D contains a derivation of the distribution of counts in randomly placed cells by extending some of the ideas developed in this paper. Although all the results of this paper are independent of those derived in Appendix D, it has been included because the final result in it (equation D3) is known to be accurate. Thus, Appendix D provides one natural way in which the usual Press–Schechter analysis may be extended to provide additional information about the evolution of nonlinear gravitational clustering.

2 A RELATION BETWEEN THE SPREAD OF DISEASE AND THE PRESS–SCHECHTER APPROACH

Consider a disease that is spread in accordance with the following model. Assume that, initially, there is a single carrier of the disease. Assume that this carrier is capable of infecting others, and that the probability that it infects k others is given by a Poisson distribution. If the initial carrier is thought of as belonging to the zeroth generation, then each of these newly infected carriers belongs to the first generation. Assume that each of the members of the first generation is, in turn, capable of infecting still others, who will make up the second generation. The members of the second generation infect still others who make up the third generation, who infect still others, and so on. Assume that, in any generation, the probability that a carrier is able to infect k others is given by a Poisson distribution that is specified by the parameter, b , say. It is possible that, by chance, none of the carriers in the n th generation infects any new members. In this case the number of members in the $(n+1)$ th generation is zero. If this should happen, the spread of the disease will be said to be halted, and we can ask for the probability that there were N people infected in total, including the initial member in the zeroth generation.

This model for the spread of disease has been studied in some detail. It is a Galton–Watson branching process in which the distribution of the number of progeny of a given member of each generation is a Poisson distribution with mean $b < 1$ (see, e.g., Harris 1963). This Galton–Watson process has an analytic solution. The probability that N people were infected in total is given by a Borel distribution with parameter b (Otter 1949; Good 1960; Consul 1989).

It is possible to use this Poisson Galton–Watson branching process to study the growth of gravitational clustering from an initially Poisson distribution. Consider a randomly chosen point in a Poisson distribution; this point comprises the zeroth generation. All points that are within a given ‘contagious’ volume, say v_c , around this point will be considered to be infected. Since the distribution is Poisson, the




Figure 1. A clump with six member particles identified in the initial particle distribution using the friends-of-friends percolation model. In the percolation model, clumps merge with each other as the link length used to define friends-of-friends increases. The overlapping of volumes makes the percolation model difficult to treat analytically.

probability that this point is able to infect k others is Poisson, and the parameter that specifies this Poisson distribution is related to the distance from the carrier out to which the disease is contagious. For convenience, assume that v_c is defined so that the parameter of the Poisson distribution is $b = \bar{n}v_c$ where $\bar{n}$ is the average density. These k points make up the members of the first generation. However, each member of the first generation will also have been able to ‘infect’, say, j others, corresponding to the j neighbours that could have been (Poisson distributed) within v_c from it. The set of all these neighbours of all the k members of the first generation makes up the second generation, and so on. It is clear that this situation is similar to the one described in the previous paragraph. This means that this Galton–Watson model for clustering from an initially Poisson distribution implies that the distribution of nonlinear clump sizes is a Borel distribution. In other words, this Galton–Watson model provides the same description of nonlinear clustering as the better known Press–Schechter type analyses described earlier.

For this description to be exactly like the Galton–Watson process described above, we must assume that the probability that one of the members in the first generation has j neighbours within v_c from it is independent of the fact that it is one of k particles within v_c from the initial particle in the zeroth generation. This is the same as assuming that the volume v_c centered on the initial particle does not even partially overlap the volume v_c centered on each of the k members in the first generation. Clearly, this assumption is false; the assumption that the volumes never overlap is a gross simplification. Nevertheless, we will continue considering this simplified model, since it provides the same mass function as the Press–Schechter Borel distribution.

One can argue that the simplification which enabled us to pose the problem in terms of this Galton–Watson model suggests one way in which the Press–Schechter mass functions could be modified. Accounting for the fact that it is possible for the volume v_c centered on the initial particle to partially overlap the volume v_c centered on one of the k members in the first generation, and so on, is the problem known as friends-of-friends percolation. Clearly, percolation




Figure 2. A branching process model of hierarchical clustering. The number of particles in each oval is a random variable determined by the (Poisson) initial conditions. The two ovals for each particle represent two different epochs, corresponding to two different link lengths, or two different over-density thresholds, with oval size increasing (over-density threshold decreasing) with time. In this example, the clump of six particles was composed of four single particles and one pair at the earlier epoch. The fact that different volumes are assumed to not overlap makes this branching process analytically tractable.

is distinct from the Galton–Watson process described above. The percolation model as formulated here can be developed as an alternative model for the growth of clustering. In principle, friends-of-friends clump mass functions can be calculated from the initial distribution; they are functions of the percolation link length (which is simply related to the size of the ‘contagious’ volume v_c), and are different from the clump mass functions determined using the Galton–Watson model. Moreover, mergers are also well-defined in the percolation model, as is the partition function of merger trees. Thus, the percolation model is at least as well defined as the Press–Schechter excursion set model. However, at present, neither the distribution of friends-of-friends clump sizes, nor the associated merger probabilities can be calculated analytically. Since the Galton–Watson model provides the same distribution of clump masses as do the excursion set, or the cloud-in-cloud, analyses of the Press–Schechter approach, the percolation model will not be considered further in this paper.

3 THE GALTON–WATSON DESCRIPTION OF HIERARCHICAL CLUSTERING

The excursion set formulation of the Press–Schechter theory shows clearly how to formulate and solve for a description of merging and hierarchical clustering (Bond *et al.* 1991; Lacey & Cole 1993). Essentially, the merging problem reduces to solving a two-barrier problem that is analogous to the one-barrier level crossing problem that was solved to obtain the Press–Schechter multiplicity function. For an initially Poisson distribution the two-barrier problem has also been solved. The probability that, at the epoch b_1 , one of the progenitors of a clump with k particles at the epoch $b_2 > b_1$, was of size $j \leq k$, is easily related to the conditional distribution

$$f(j, b_1 | k, b_2) = \binom{k}{j} \frac{j^j}{k^k} \left(\frac{b_1}{b_2} \right)^{j-1} k \left(\frac{b_2 - b_1}{b_2} \right) \times \left[k \left(\frac{b_2 - b_1}{b_2} \right) + (k - j) \frac{b_1}{b_2} \right]^{k-j-1} \quad (4)$$

where $b_1 < b_2$ and $k \geq j$ (equation 40 in Sheth 1995). Here, $f(j, b_1 | k, b_2)$ is the probability that a particle which is in a k -particle clump at the epoch b_2 was in a j -particle clump at the epoch b_1 . The statements $f(j, b_1 | k, b_2)$ are independent of what happened at times previous to b_1 and also of what will happen at times later than b_2 . Appendix B shows that, in the appropriate limits (i.e., $k \gg 1$, $j \gg 1$, and $k - j \gg 1$, and also $b_1 \rightarrow 1$ and $b_2 \rightarrow 1$), equation (4) reduces to equation (2.15) of Lacey & Cole (1993) (also see Fig. 2 and associated algebra in Sheth 1995).

Following Lacey & Cole (1993), equation (4) can be manipulated to provide an expression for the probability that a clump with k particles at the epoch b_2 was formed from other subclumps that merged with a subclump that had exactly j particles at the epoch b_1 . However, it does not provide any information about the distribution of sizes of the other subclumps, other than the restriction that the total number of particles in those other subclumps must sum to $k - j$. Another way of thinking of $f(j, b_1 | k, b_2)$ is to note that it is obtained without consideration of the different ways in which the object of size j (at the epoch b_1) could have merged with other objects to form the final object of size k (at the epoch b_2). If the merging process is visualized as a tree (so trees having different numbers of branches, branch sizes and branching points describe different merger histories), a useful quantity is the probability that a particular tree structure, rather than any other, occurs. So, the problem is to solve for what is, essentially, the partition function for various tree structures.

Recall that the statements $f(j, b_1 | k, b_2)$ are independent of what happened at times previous to b_1 and also of what will happen at times later than b_2 . So, to solve for the entire merger history tree (i.e., for many $b_1 < b_2 < \dots$) one need only know how to solve for the tree structure at any given two epochs, say, b_1 and b_2 . Therefore, in what follows the later epoch, b_2 , is referred to as the final epoch. Notice also that the statements $f(j, b_1 | k, b_2)$ imply that it is possible to calculate the partition function for a tree with k particles without explicitly considering the partition function for trees with $l \neq k$ particles. So, for any given pair of epochs b_1 and b_2 , and for any tree whose final size (i.e., at the epoch b_2) is k , the problem is to calculate all possible tree configurations, and to assign to each configuration the probability that it actually occurs. The elements of the set of all possible k -particle tree configurations are simply the various ways of partitioning the integer k (physically, this is what is required by mass conservation).

Let $p(1^{n_1} 2^{n_2} \dots k^{n_k} | k)$, with $n_1 + \dots + n_k = m$ and $\sum_{j=1}^k j n_j = k$, denote the probability that the final clump with k particles at b_2 had m progenitors at b_1 , of which n_1 were singles, n_2 were pairs, n_j were subclumps with j particles each, and so on. Note that $p(1^{n_1} 2^{n_2} \dots k^{n_k} | k)$ is independent of the order of $\{1^{n_1} 2^{n_2} \dots k^{n_k}\}$ since different permutations of the ‘branches’ should all have the same probability of occurring. The problem is to calculate this probability for given k, m, b_2, b_1 and $\{n_1, n_2, \dots, n_k\}$. To date, it has not proven possible to calculate this probability exactly from the excursion set description itself (Kauffmann & White 1993; Sheth 1995). In this section we will use the Galton–Watson interpretation of the clustering process to obtain probabilities for the various partitions of k .

The Galton–Watson process can be used to formulate a

description of the merger history of a clump as follows. As before, consider the single member of the zeroth generation. This person has a number of children, and they each have some number of children, and so on down the family tree. If the probability a given member of the tree has n children is given by a Poisson distribution with parameter b_2 , then we have the same branching process as before. The probability that such a family tree has N members in total, after which the family died out, is given by a Borel distribution with parameter b_2 . Now consider the more complicated case in which some of the children are male and some female. Assume that the probability that a given member on the tree has n_1 daughters is a Poisson distribution with parameter b_1 and the probability of having n_2 sons is a Poisson distribution with parameter $b_2 - b_1$. This is the same as assuming that the probability that a given member on the tree has n children is given by a Poisson process with parameter b_2 , and the probability that n_1 are daughters and $n_2 = n - n_1$ are sons is given by a Binomial distribution where the probability that any given child is female is $p = b_1/b_2$. Consider the family tree of such a process, and assume that the initial ancestor was male. For such a family tree, we can ask for the probability that there were N_1 females and $N_2 = N - N_1$ males in total in the family tree. This branching process is a special case of that studied by Good (1960).

Now imagine drawing the family tree. Use dots to represent children of either sex, but only draw lines between parents and their daughters. Then the tree will consist of a number of groups, in which each member of the group is connected to other members of the same group, but not to members from different groups. Call these groups subfamilies. (The tree is inherently sexist, as the first member at the head of every new subfamily is always male.)

We will characterize trees by the distribution of the sizes of the subfamilies in them. Let n_i denote the number of subfamilies each with i members. Then we can ask for the probability $p(1^{n_1} \dots k^{n_k} | k)$ that a tree with k members has exactly m subfamilies (so $n_1 + \dots + n_k = m$), and that there were n_1 singles, n_2 pairs, n_j j -tuples, and so on. Clearly, this characterization of a family tree (by the size and number of subfamilies in it) is exactly analogous to the distribution of subclump sizes for a given Press–Schechter clump. It provides us with probabilities for the various partitions of k . In particular, this extension of the Poisson Galton–Watson branching process requires that

$$p(1^{n_1} 2^{n_2} \dots k^{n_k} | k) = \frac{\eta(1, b_1)^{n_1} \eta(2, b_1)^{n_2} \dots \eta(k, b_1)^{n_k}}{\eta(k, b_2)} \times \frac{m!}{n_1! n_2! \dots n_k!} \times \frac{1}{m} \frac{[k(b_2 - b_1)]^{m-1} e^{-k(b_2 - b_1)}}{(m-1)!} \quad (5)$$

where $\eta(l, b)$ is the Borel distribution with parameter b (eq. 1), and n_j denotes the number of subfamilies in the tree that have exactly j members (with the usual convention that $0! = 1$). Since this branching process is related to the gravitational clustering process, the probability that a clump of size k at the epoch b_2 had the m progenitors $n_1, n_2, \dots, n_k$ at the epoch b_1 is given by equation (5). Thus, equation (5) can be used to generate the partition function of merger history trees. It is the main result of this paper.

Appendix B shows that this partition function is consistent with the merger probabilities of equation (4). It also provides additional insight into the origin of the various terms in equation (5). Appendix C describes a queueing process that provides another way to derive this expression.

3.1 Some properties of the partition structure

Some limits of equation (5) are worth studying. As $b_2 \rightarrow b_1$, $p(1^{n_1} \dots k^{n_k} | k) \rightarrow 0$, except when $m = 1$, for which $p(k^1 | k) \rightarrow 1$ as required. As $b_1 \rightarrow 0$, then $\eta(l, b_1) \rightarrow 0$ except for $l = 1$, so that

$$\begin{aligned} \lim_{b_1 \rightarrow 0} p(1^{n_1} \dots k^{n_k} | k) &= p(1^k | k) \\ &= \frac{(kb_2)^{k-1} e^{-kb_2}}{k!} \frac{1}{\eta(k, b_2)} = 1. \end{aligned} \quad (6)$$

In other words, in the limit as $b_1 \rightarrow 0$, all progenitors are certainly single particles, which is also the expected result.

The probability $n(m|k)$ that a clump of size k at the epoch b_2 has m progenitors at the epoch b_1 is given by summing $p(1^{n_1} \dots k^{n_k} | k)$ over all distinct sets of m integers that add up to k . Now, any set specified by $\{n_1, \dots, n_k\}$ with $n_1 + \dots + n_k = m$ can be written explicitly in terms of its m members as $\{l_1, \dots, l_m\}$. Thus, $l_1 + \dots + l_m = \sum_{j=1}^m j n_j = k$. So,

$$\begin{aligned} n(m|k) &= \sum_{m \text{ parts.}} \frac{\eta(1, b_1)^{n_1} \dots \eta(k, b_1)^{n_k}}{\eta(k, b_2)} \frac{m!}{n_1! \dots n_k!} \\ &\quad \times \frac{1}{m} \frac{[k(b_2 - b_1)]^{m-1} e^{-k(b_2 - b_1)}}{(m-1)!} \\ &= \sum_{m \text{ parts.}} \left(\frac{1^{1-1}}{1!} \right)^{n_1} \dots \left(\frac{k^{k-1}}{k!} \right)^{n_k} \\ &\quad \times \frac{b_1^{1n_1 + \dots + kn_k - n_1 - \dots - n_k}}{e^{1n_1 b_1 + \dots + kn_k b_1}} \\ &\quad \times \frac{k!}{(kb_2)^{k-1} e^{-kb_2}} \times \frac{m!}{n_1! \dots n_k!} \\ &\quad \times \frac{1}{m} \frac{[k(b_2 - b_1)]^{m-1} e^{-k(b_2 - b_1)}}{(m-1)!} \\ &= \frac{k!}{m!} \frac{1}{k^{k-m}} \left(\frac{b_1}{b_2} \right)^{k-m} \left(1 - \frac{b_1}{b_2} \right)^{m-1} \\ &\quad \times \sum_{\text{dist. perms.}} \frac{l_1^{l_1-1} l_2^{l_2-1} \dots l_m^{l_m-1}}{l_1! l_2! \dots l_m!} \\ &= \frac{k!}{m!} \frac{1}{k^{k-m}} \left(\frac{b_1}{b_2} \right)^{k-m} \left(1 - \frac{b_1}{b_2} \right)^{m-1} \frac{m}{k} \frac{k^{k-m}}{(k-m)!} \\ &= \binom{k-1}{m-1} \left(\frac{b_1}{b_2} \right)^{k-1-(m-1)} \left(1 - \frac{b_1}{b_2} \right)^{m-1}. \end{aligned} \quad (7)$$

The sum in the third equality is over all sets of m integers which satisfy $l_1 + \dots + l_m = k$, and over all the distinct permutations of each set (which accounts for the multinomial factor $m!/n_1! \dots n_k!$). The second from last equality follows from a combinatorial identity (note the similarity to the Borel–Tanner distribution of equation B3; also see Sheth & Saslaw 1994). The final expression for $n(m|k)$ is the same as that obtained previously (eq. 52 in Sheth 1995).

It is easy to see that this expression is sensible. Referring back to the Galton–Watson family tree description, recall

that the head of the tree is always the initial male, and other males start subfamilies within the tree. So, the number, m , of subfamilies within a family tree having k members in total is equal to the number of male members in the tree. Now, the probability any given child is male is $q = 1 - p = (b_2 - b_1)/b_2$. Since it is certain that one member of the tree is a male (the head of the family tree is always a male), the probability that there are m males in a family with k members should be $[(k-1)!/(k-m)!(m-1)!] p^{k-m} q^{m-1}$. This is identical to the Binomial distribution of equation (7). Alternatively, this expression for the number of subclumps of a k -sized clump, which is the same as the number of sons in a k sized family, can be obtained directly from Good’s (1960) treatment of the Galton–Watson process with many types of progeny. The brute force calculation is given in Appendix A. It, too, shows that $n(m|k)$ is given by equation (7).

Equation (7) is also consistent with the physical requirement that, in the limit of very small time steps, the probability that a clump has two progenitors should be an infinitesimal, the probability that the clump has three progenitors should be an infinitesimal of the next higher order, and so on. It also implies that massive clumps form preferentially from massive progenitor clumps: on average, the size, at some earlier epoch, of the largest progenitor clump of a clump at some given final epoch depends significantly on how massive the final clump is relative to the characteristic mass at the final epoch (eq. 55 in Sheth 1995).

In the limit of large k and large subclumps l_i , equation (5) has an interesting interpretation. Stirling’s formula for the factorials implies that

$$\begin{aligned} p(l_1, \dots, l_m | k) &= \frac{l_1^{l_1-1} \dots l_m^{l_m-1}}{l_1! \dots l_m!} \frac{k!}{k^{k-1}} \frac{k^{m-1}}{n_1! \dots n_k!} \\ &\quad \times \left(\frac{b_1}{b_2} \right)^{k-m} \left(1 - \frac{b_1}{b_2} \right)^{m-1} \\ &\rightarrow \frac{k^{m-1}}{n_1! \dots n_k!} \left(\frac{b_1}{b_2} \right)^{k-m} \left(1 - \frac{b_1}{b_2} \right)^{m-1} \\ &\quad \times \frac{\sqrt{2\pi} k^{\frac{3}{2}}}{(2\pi)^{\frac{m}{2}}} \prod_{i=1}^m \frac{1}{l_i^{3/2}} \\ &\rightarrow \frac{1}{n_1! \dots n_k!} \left(\frac{b_1}{b_2} \right)^{k-m} \left(1 - \frac{b_1}{b_2} \right)^{m-1} \\ &\quad \times \frac{1}{(2\pi k)^{(m-1)/2}} \prod_{i=1}^m \frac{1}{x_i^{3/2}}, \end{aligned} \quad (8)$$

where we have defined $x_i \equiv l_i/k$. The first term on the right is the combinatorial term that is included to insure that each distinct combination of subclumps is counted only once. The second set of terms accounts for the probability that there are exactly m subclumps. The final product term is the most interesting. It shows that, for a given value of the final clump size k , and if $m \ll k$, the case in which the subclumps are all approximately the same size is much less likely than the case in which one of the subclumps is very much more massive than all the others.

The partition function (eq. 5) exhibits an interesting scaling. It depends on the initial and final epochs, b_1 and b_2 , only through the combination $p = b_1/b_2$. As one would expect, this is also true of the simpler statement given in equation (4). This means that the probability $p(l_1, \dots, l_m | k)$, given the two epochs b_1 and b_2 , will be the same as the

probability $p'(l_1 \dots, l_m | k)$, given b'_1 and b'_2 , provided that $b_1/b_2 = b'_1/b'_2 = p$. In this sense, the clustering evolves in a self-similar fashion. This scaling can be exploited when comparing equation (5) with merger histories of clumps in N -body simulations. When $b_1 \rightarrow 1$ and $b_2 \rightarrow 1$, the ratio $b_1/b_2 = (1 + \delta_{c2})/(1 + \delta_{c1}) \rightarrow 1 + \delta_{c2} - \delta_{c1}$. Thus, the scaling in b_1/b_2 corresponds to the scaling in $(\delta_{c1} - \delta_{c2})$ discussed, for example, by Bond *et al.* (1991) and by Lacey & Cole (1993).

4 APPLICATIONS

One example of the type of statistical question that we can now answer is as follows. Suppose we are interested in the distribution of sizes of the largest progenitor clump at some epoch b_1 , of a given clump at the epoch b_2 . This might be of interest, for example, in studies of the Butcher–Oemler effect (Bower 1991; Kauffmann 1995). It is straightforward to compute the relevant sums over the partition function numerically. As an example, Fig. 3 shows the probability that the largest subclump of a clump with k particles has l_1 particles, for two choices of k , and for two choices of the ratio $p = b_1/b_2$ of the initial and final epochs.

Three features of Fig. 3 are obvious. First, for given k , the curves depend strongly on p , the ratio of the two epochs b_1 and b_2 . As p increases, the curves peak at higher values of l_1/k . This means that the largest progenitor of a given clump is a smaller fraction of the total mass as the time between the final epoch and the epoch at which the progenitors were identified increases. This simply reflects the fact that the clustering is hierarchical; small clumps merge to form big clumps, and on average, clumps were smaller in the more distant past than they are at present. Second, the curves depend strongly on k , the number of particles in the final clump. For a given value of p , curves with higher values of k peak further towards the left. That is, for a given value of p , the largest progenitor of a massive clump is more likely to be a smaller fraction of the total mass than is the largest progenitor of a less massive clump. In this sense, for any choice of initial and final epochs (because the partition function only depends on the ratio b_1/b_2), more massive clumps always appear to have assembled a larger fraction of their mass more recently than less massive clumps. Lacey & Cole (1993) show that this is also what happens in the Gaussian case. Furthermore, it is consistent with the analytical result that, for a clump that has k particles in total, the average number of particles in a progenitor subclump (not necessarily the largest subclump) is $k/[p + k(1 - p)]$ (Sheth 1995).

So, we expect the average size, and the most probable size, of the largest progenitor clump to decrease as p decreases. However, based on Fig. 3, the scaling property of the partition function, and the fact that b evolves quickly initially and slower at later times (see discussion following eq. 2), we can also study the rate with which this size decreases. As a specific example, consider two clumps, each of size of k . Assume that one of the clumps is completely assembled at epoch b_2 and the other at $b'_2 < b_2$ (the primed clump was assembled earlier than the unprimed clump). Now consider the progenitors of the unprimed clump that are identified at the epoch $b_1 = pb_2$. The partition function (eq. 5)




Figure 3. Probability, $f(l_1|k)$, that the largest subclump of a k particle clump has l_1 particles, for two choices of the ratio $p = b_1/b_2$ of the initial and final epochs, and for two choices of k . Solid lines show the distribution for $k = 10$ with $p = 0.3$ and 0.9 . Dashed lines show the distribution for $k = 50$, with the same values of p .

specifies this distribution of progenitor subclumps. So, we can compute, e.g., the average size of the largest progenitor subclump. Now consider the primed clump. The scaling property of the partition function shows that its subclump distribution will be the same as that of the unprimed clump at the epoch when $b'_1 = pb'_2 < b_1$, where $p = b_1/b_2$ has the same value as for the unprimed clump. Consider the average size of the largest progenitor of these two clumps as a function of ‘lookback time’ from the epochs b_2 and b'_2 .

Since p is the same for the two clumps, they will have the same average size for the largest subclump at the epochs b_1 and $b'_1 < b_1$. However, the lookback time for the unprimed and the primed clumps is the time corresponding to $b_{21} = b_2 - b_1$ and $b'_{21} = b'_2 - b'_1$, respectively. Since b changes ever more slowly as it increases, $b'_{21} < b_{21}$. This implies that the average size of the largest progenitor of the primed clump (which was assembled earlier) decreases more rapidly than it does for the unprimed clump (which was assembled later). In other words, when phrased in terms of lookback time, the evolutionary history of a clump of size k depends on when it was first assembled.

As discussed above, the evolutionary histories of clumps, when phrased in terms of lookback time, depend on the rate of change of b . In the previous paragraph we were able to draw conclusions about the dependence of evolution on formation epoch because the rate of change of b is a function of epoch. However, as noted in the introduction, the rate of change of b also depends on the background cosmology. Thus, one also expects the evolutionary histories of clumps to be sensitive to the background cosmology. For example, the $b(t)$ curves for different cosmologies show that a clump of a given mass will have formed at a greater lookback time in a low density universe than in one with critical density. These trends, the more recent assembly of larger relative to smaller clumps, and the sensitivity of merger histories to the background cosmology, have been noted by Lacey & Cole (1993) and by Kauffmann (1995) in their study of clustering from initially Gaussian fields. The implication that similar mass clumps which were assembled at different times have different lookback time histories is in qualitative agreement

with the numerical, Monte–Carlo model used by Kauffmann (1995).

The third feature that is evident in Fig. 3 is simply that the curves are extremely skew. This means that the average number of particles in the largest progenitor subclump is not necessarily a good indicator of the most probable number of particles in the largest progenitor. Therefore, the curves of average merger histories given in Fig. 1 of Kauffmann (1995) should be treated carefully. Note that equation (5) provides an efficient way of evaluating the difference between the mean and the most probable sizes.

In addition to allowing one to calculate the dispersion around the mean history of any given clump, the partition function can also be used to compare this Galton–Watson Poisson Press–Schechter model with N -body simulations. It is also useful to compare the merger histories described by equation (5) with the ad hoc, Monte–Carlo merger histories generated by Kauffmann & White (1993). This will provide a test of the Galton–Watson model developed here, and may also provide some insight into the reason for the accuracy of the Kauffmann–White algorithm.

To effect this comparison, we will plot the ratio of the largest progenitor clump to the total mass l_1/k , versus the ratio of the second largest progenitor to the largest progenitor, l_2/l_1 . Figures 2 and 3 in Kauffmann & White (1993) show examples of such plots. Fig. 4 (of this paper) shows what is essentially the joint probability that the largest two progenitor subclumps are l_1 and l_2 , given that the final clump is of size k , as determined by the partition function (eq. 5) for $p = b_1/b_2 = 0.8$ (top panel) and 0.3 (bottom panel), and for $k = 10$ (solid contours) and $k = 50$ (dotted contours). Large values of p correspond to large time differences between the initial and final epochs. The results are plotted in terms of the ratio l_1/k versus l_2/l_1 , for ease of comparison with the work of Kauffmann & White (1993). The contours are at 0.0001, 0.0003, 0.001, 0.003, 0.01, 0.03, and 0.1. The figure shows that, for a given value of k , most of the clumps lie along a relatively narrow band in the $(l_1/k), (l_2/l_1)$ plane. The location of the band depends on the values of k and p .

Fig. 4 is qualitatively similar to Figs. 2 and 3 in Kauffmann & White (1993). At small lookback times (large p , top panel) the band lies at somewhat larger values of l_1/k than at larger lookback times (small p , bottom panel). This simply reflects the fact that, at smaller lookback times, a larger fraction of the clump survives relatively intact. However, the band for the clump with larger k is at a lower value of l_1/k than the less massive clump. This is consistent with the faster assembly required by larger clumps that we deduced in Fig. 3. The top panel also shows that, in this (relatively small lookback time) regime, the joint probability distribution has one peak at large values of l_1/k (and correspondingly small values of l_2/l_1), and a broader, not so high peak around the values of $l_1 \sim k/2$ and $l_2 \sim l_1/2$. Thus, at small lookback times, it appears that most clumps grow because a large clump accretes many smaller ones. Those that do not, grow because of the mergers of objects that are approximately the same size.

At larger lookback times (bottom panel), the band is shifted towards lower values of l_1/k . Furthermore, the joint probability distribution becomes peaked towards the lower right hand corner of the plot, at values of $l_2/l_1 \approx 1$. This,




Figure 4. Joint probability that the largest two progenitor subclumps are l_1 and l_2 , given that the final clump is of size k as determined by the partition function (eq. 5) for $p = b_1/b_2 = 0.8$ (top panel) and $p = 0.3$ (bottom panel), when $k = 10$ (solid contours) and $k = 50$ (dotted contours). The results are plotted in terms of the ratio l_1/k versus l_2/l_1 , for ease of comparison with the earlier work described in the text. The contours are at probability levels of 0.0001, 0.0003, 0.001, 0.003, 0.01, 0.03, and 0.1.

too, is sensible, since at large lookback times, most clumps have split up into a large number of small, approximately equally massive progenitors. These features are in qualitative agreement with those measured in N -body simulations (Kauffmann & White 1993, and references therein).

5 DISCUSSION

The partition function that describes the relative probabilities of all possible merger histories of those Press–Schechter clumps which form as an initially Poisson distribution evolves gravitationally, can be written in closed form (eq. 5). The partition function is a function of the initial and final epochs (denoted by b_1 and b_2 , respectively). Since the time evolution of b can be computed (cf. the Introduction) the temporal evolution of the partition function is known. The counts in cells distribution associated with this Poisson Press–Schechter distribution can also be computed by extending the branching process analogy (Appendix D). It, too, is a function of b , so its temporal evolution is also

known. Thus, the Press–Schechter description of clustering from an initially Poisson distribution is now complete.

It may be worth pointing out that Appendix B shows that the Poisson Galton–Watson branching process specifies the Borel clump size distribution (equation 1) and the merger probabilities (equation 4) uniquely, and also allows one to describe the merger tree completely (equation 5). On the other hand, if equations (1) and (4) are treated as the only known constraints on the form of the merger tree, then the branching process considered in this paper is not the only way to construct the merger history trees consistent with these constraints (e.g., Sheth 1995). For instance, one can always construct an ad hoc Monte-Carlo scheme, like that of Kauffmann & White (1993), which satisfies equations (1) and (4), and is not necessarily consistent with the branching process description of equation (5). For this reason we have argued that, rather than being completely ad hoc, the branching process is, indeed, a reasonable model for the growth of clustering (Figs. 1 and 2 and associated discussion).

A Poisson distribution is similar to a white noise Gaussian random field. So, this result can be related to the evolution of clustering from an initially Gaussian density field that has a scale free power spectrum with slope $n = 0$. Can it be extended to describe the growth of gravitational clustering from Gaussian random fields with arbitrary initial power spectra?

One way in which to do this is as follows. The Borel distribution reduces to the Press–Schechter multiplicity function in the limit as $N \gg 1$ and as $b \rightarrow 1$. Similarly, $f(j|k)$ (eq. 4) reduces to the $n = 0$ Gaussian result provided that $k \gg 1$, $j \gg 1$, $k-j \gg 1$, and b_1 and $b_2 \rightarrow 1$ (Sheth 1995). For a given density threshold δ_c [eq. 2 shows that $b = 1/(1+\delta_c)$], Gaussian fields with different power spectra yield different excursion set Press–Schechter mass multiplicity functions. However, these differences arise solely because the relation between the variance and the mass depends on the power spectrum. When written directly in terms of the variance, rather than the mass, the Press–Schechter multiplicity functions and the merger probabilities have a universal form that is independent of the underlying power spectrum (e.g. Lacey & Cole 1993). This suggests writing the partition function of equation (5) in terms of the variance of the Poisson ($n = 0$) distribution. Then, in the large l_i and $b \rightarrow 1$ limits, equation (5) should reduce to the Gaussian result. The appropriate Jacobian can then transform the partition function (written in terms of the variance) into an expression for the masses of the subclumps.

There may be a more sophisticated way to describe the merger histories of clumps that form from Gaussian random fields. In this paper, branching processes and queueing theory were both used to formulate and solve a problem that was originally posed in terms of trees associated with random walk excursions on a discrete Poisson distribution. So the question is, Is it possible to extend any of these relations to the continuous Gaussian case? For example, is it possible to formulate the excursion sets of random walks associated with Gaussian fields, described by Bond *et al.* (1991) in their derivation of the Press–Schechter mass functions, in terms of branching processes? Trees associated with excursions of random walks on Gaussian fields are the subject of current interest (Neveu & Pitman 1989; Le Gall 1993). The $b \rightarrow 1$

limit is known as the ‘critical’ Galton–Watson process (the $b < 1$ case considered in this paper is ‘subcritical’). Many properties of the tree associated with this critical process have been calculated (Aldous 1993 and references therein). The application of these ideas to the Press–Schechter description of clustering from arbitrary Gaussian initial conditions is in progress.

One final way in which to extend the ideas of this paper to arbitrary initial conditions is to note that the usual Press–Schechter mass functions provide a good, but by no means perfect, description of N -body simulations of clustering. Thus, one might reconsider the percolation model discussed in Section 2. First, one would compare the percolation mass functions (obtained numerically) for the Poisson case with those of the branching process (the Borel distribution, equation 1). If the percolation mass functions provide an acceptable fit to the simulation results, then one could also compute (numerically) and test the percolation merger probabilities and merger tree. Of course, it is trivial to apply the percolation model to arbitrary initial conditions.

The discussion at the end of Section 2 shows clearly that this branching process extension of the Press–Schechter approach is correct only in a strictly statistical sense. This is because, unlike in the percolation model discussed in Section 2, there is no direct correspondence between particles in the initial distribution and particles in clumps. That is, the Galton–Watson branching process, like the excursion set interpretation of the Press–Schechter mass function, is a purely statistical model for the growth of clustering. To be useful, it relies on the accuracy and applicability of the ergodic hypothesis. It cannot, and should not, be expected to work on a particle-by-particle basis.

ACKNOWLEDGMENTS

I thank I. M. and R. G. Hook for much needed encouragement, G. Lemson for arranging a trip to the island of Schiermonnikoog where my interest in branching process descriptions of clustering began, D. Aldous for stimulating discussions, J. Pitman for providing valuable insight and entertaining cricket, and the referee, C. Lacey, for suggestions that improved the presentation of these results.

REFERENCES

- Aldous D. J., 1993, *Ann. Prob.*, 21, 248
- Bond J. R., Cole S., Efstathiou G., Kaiser N., 1991, *ApJ*, 379, 440
- Borel E., 1942, *Compt. rend.*, 214, 452
- Consul P. C., 1989, *Generalized Poisson Distributions: Properties and Applications*. M. Dekker, New York
- Efstathiou G., Frenk C., White S. D. M., Davis M., 1988, *MNRAS*, 235, 715
- Epstein R. I., 1983, *MNRAS*, 205, 207
- Good I. J., 1960, *Proc. Camb. Phil. Soc.*, 56, 367
- Harris T. E., 1963, *The Theory of Branching Processes*. Springer-Verlag, Berlin
- Jedamzik K., 1995, *ApJ*, 448, 1
- Kauffmann G., 1995, *MNRAS*, 274, 200
- Kauffmann G., White S. D. M., 1993, *MNRAS*, 261, 921
- Kendall D. G., 1951, *J. Roy. Stat. Soc. B*, 13, 151
- Lacey C., Cole S., 1993, *MNRAS*, 262, 627

- Lacey C., Cole S., 1994, MNRAS, 271, 676
 LeGall J. F., 1993, Probab. Th. Rel. Fields, 96, 369
 Neveu J., Pitman J., 1989, Lecture Notes Math, 1372, 248,
 Séminaire de Probabilités XXIII. Springer–Verlag, Berlin
 Otter R., 1948, Ann. Math. Stat., 20, 206
 Press W., Schechter P., 1974, ApJ, 187, 425
 Riordan J., 1979, Combinatorial Identities. Robert E. Krieger
 Publ. Co., New York
 Saslaw W. C., 1989, ApJ, 341, 588
 Saslaw W. C., Hamilton A. J. S., 1984, ApJ, 276, 13
 Sheth R. K., 1995, MNRAS, 276, 796
 Sheth R. K., Saslaw W. C., 1994, ApJ, 437, 35
 Tanner J. C., 1953, Biometrika, 40, 58
 Tanner J. C., 1961, Biometrika, 48, 222

APPENDIX A: THE POISSON GALTON–WATSON PROCESS WITH TWO TYPES OF PROGENY

The Poisson Galton–Watson process with m different types of progeny has been studied by Good (1960). He shows that if the branching process starts with r_i individuals of type i , for all $1 \leq i \leq m$, then the probability that the whole tree contains exactly n_1 of type 1, n_2 of type 2, and so on, is equal to the coefficient of

$$z_1^{n_1-r_1} z_2^{n_2-r_2} \dots z_m^{n_m-r_m}$$

in

$$f_1^{n_1} \dots f_m^{n_m} \left\| \delta_\mu^\nu - \frac{z_\mu}{f_\mu} \frac{\partial f_\mu}{\partial z_\nu} \right\|,$$

where δ_μ^ν is the identity matrix and

$$f_\mu(z) = \prod_\nu \exp(-a_{\mu\nu}(1-z_\nu))$$

reflects the fact that the probability that an individual of type μ has a child of type ν is a Poisson distribution with parameter $a_{\mu\nu}$ independently of the other individuals.

For our problem, $m = 2$, and subscript 1 is for girls and subscript 2 is for boys. Then $a_{11} = a_{21} = b_1$ and $a_{12} = a_{22} = b_2 - b_1$, and the problem is to extract the relevant coefficients of

$$\left\{ \exp\left(n_1 [b_1(z_1 - 1) + (b_2 - b_1)(z_2 - 1)] + n_2 [b_1(z_1 - 1) + (b_2 - b_1)(z_2 - 1)]\right) \right\} \times \left\{ [1 - z_1 b_1] [1 - z_2(b_2 - b_1)] - z_2 b_1 z_1(b_2 - b_1) \right\}. \quad (\text{A1})$$

Suppose that we are interested in the family tree that was started by one male ancestor, in which there are N members of the tree in total. Then we can use equation (A1) to calculate the probability of having a tree with exactly N members. Let n denote the number of females in the tree. Then set $n_1 = n$ and $n_2 = N - n$ and let $r_1 = 0$ and $r_2 = 1$. The answer is obtained by summing up all the coefficients of terms of order $z_1^n z_2^{N-n-1}$. This involves straightforward but tedious algebra which is not reproduced here. The result is

that the probability of having N members in the tree is

$$\begin{aligned} P(N) &= \eta(N, b_2) \\ &\times \sum_{n=0}^{N-1} \binom{N-1}{n} \left(\frac{b_1}{b_2}\right)^n \left(1 - \frac{b_1}{b_2}\right)^{N-1-n} \\ &= \eta(N, b_2), \end{aligned} \quad (\text{A2})$$

where n denotes the number of girls in the tree. Clearly, this is the same result as that given by equation (7) in the main text.

APPENDIX B: MERGER PROBABILITIES AND THE PARTITION FUNCTION

Some of the results of this Appendix were suggested by the recent work of Pitman (in preparation). This Appendix shows explicitly that the partition function derived in the main text, equation (5) and the merger probabilities of equation (4) are consistent with each other. Demonstration of this consistency is useful because, in the appropriate limit, the merger rates implied by equation (4) are in good agreement with N -body simulations of gravitational clustering (Lacey & Cole 1994). For completeness, this limit is derived below. In addition, an expression for the mean number of j -sized subclumps of k -sized clumps is derived directly from the partition function. The argument is generalized, at the end of this Appendix, to obtain an expression for the factorial moments of this distribution.

Lacey & Cole (1993) define a merger rate by taking the limit as $\delta_1 \rightarrow \delta_2$ in $f(k, \delta_2 | j, \delta_1)$. That is, the merger rate is

$$\begin{aligned} \frac{dP(j \rightarrow k | \delta)}{d\delta} d\delta &\equiv \lim_{\delta_1 \rightarrow \delta_2} \frac{(1 - b_2) k \eta(k, b_2) f(j, \delta_1 | k, \delta_2)}{(1 - b_1) j \eta(j, b_1)} \\ &= \lim_{\delta_1 \rightarrow \delta_2} \frac{k}{(k-j)!} \frac{\delta_2}{\delta_1} \frac{(\delta_1 - \delta_2)}{(1 + \delta_2)^2} \\ &\quad \times \left(\frac{k}{1 + \delta_2} - \frac{j}{1 + \delta_1} \right)^{k-j-1} \\ &\quad \times e^{-\frac{k}{1+\delta_2} + \frac{j}{1+\delta_1}} \\ &\rightarrow \frac{k}{\sqrt{2\pi}(k-j)^{3/2}} \frac{d\delta}{(1+\delta)} \frac{e^{\delta(k-j)/(1+\delta)}}{(1+\delta)^{k-j}} \\ &\rightarrow \frac{d\delta}{\sqrt{2\pi}} \left(\frac{k^2}{(k-j)} \right)^{3/2} \frac{e^{-\delta^2(k-j)/2}}{k^2(1+\delta)} \quad (\text{B1}) \end{aligned}$$

(Sheth 1995). The third expression on the right follows from setting $\delta_1 = \delta_2 = \delta$ and $\delta_1 - \delta_2 = d\delta$, and considering the limit when $k \gg 1$, $j \gg 1$ and $k - j \gg 1$. Stirling's approximation for the factorials simplifies the expression considerably, and the final expression follows from assuming $\delta \ll 1$. Except for the $(1 + \delta)$ term in the denominator, the final expression is the same as the expression derived by Lacey & Cole (1993). In their notation $S_1 \propto 1/j$, $S_2 \propto 1/k$, and $\omega = \delta$, since the relevant Gaussian corresponding to the Poisson is the white noise case. So, in the limit where Stirling's approximation for the factorials is valid, and when $\delta \ll 1$, equation (B1) here reduces to their equation (2.17). In this limit, the merger rate implied by equation (4) describes the simulations well (Lacey & Cole 1994).

One other limit of equation (4) is also interesting. When $k \gg j$, Stirling's approximation for $k!$ and $(k-j)!$ in equa-

tion (4) implies that

$$f(j, b_1|k, b_2) \rightarrow \left(1 - \frac{b_1}{b_2}\right) \frac{j^{j-1}}{(j-1)!} \left(\frac{b_1}{b_2}\right)^{j-1} e^{-j(b_1/b_2)}. \quad (\text{B2})$$

This shows that the probability that a randomly chosen member of a Borel clump that has exactly k members at the epoch b_2 was in a Borel clump with $j \ll k$ particles at the epoch b_1 is, to an excellent approximation, given by a Borel distribution with parameter b_1/b_2 . When $b_2 \rightarrow 1$, this means that the probability that a particle is in a clump with $j-1$ other particles is given by a Borel distribution with parameter b_1 . Since the limit $b_2 \rightarrow 1$ is equivalent to requiring $\delta_2 \rightarrow 0$, this is exactly what is required by the derivation of $f(j|k)$ from the two barrier (excursion set) problem considered in Sheth (1995). Another application of Stirling's approximation (to the $j!$ term), with the limits $b_1 \rightarrow 1$ and $b_2 \rightarrow 1$ shows that equation (4) is similar to the Lacey & Cole (1993) expression for the white noise Gaussian case. Fig. 2 in Sheth (1995) also shows this to be true.

Before deriving the merger probabilities $f(j|k)$ of equation (4) directly from the partition function (equation 5), it is useful to consider some combinatorial identities. First, consider the random variable X , and assume that the distribution of X is Borel with parameter b . Then the distribution of $S_m = X_1 + \dots + X_m$, where the X_i are independent random variables, each drawn from a Borel distribution with parameter b , is given by the Borel-Tanner distribution. That is, the probability that $S_m = k$ (i.e., the sum of m independent Borel variables equals k) is

$$P(b, S_m = k) = \frac{m}{k} \frac{(kb)^{k-m} e^{-kb}}{(k-m)!}, \quad \text{where } k \geq m \quad (\text{B3})$$

(Tanner 1953, 1961). It is easy to see that this expression is sensible, since

$$\begin{aligned} & \sum_{j=1}^{k-m+1} \eta(j, b) P(b, S_{m-1} = k-j) \\ &= \sum_{j=1}^{k-m+1} \frac{(jb)^{j-1} e^{-jb}}{j!} \frac{m-1}{k-j} \frac{[(k-j)b]^{k-m} e^{-(k-j)b}}{(k-j-m+1)!} \\ &= (m-1) \frac{b^{k-m} e^{-kb}}{(k-m)!} \\ & \quad \times \sum_{i=0}^{k-m} \binom{k-m}{i} (1+i)^{i-1} (k-1-i)^{k-m-i-1} \\ &= P(b, S_m = k), \end{aligned} \quad (\text{B4})$$

as it should. The final expression follows from Abel's generalization of the Binomial theorem (see equations 14 and 20 in section 1.5 of Riordan 1979 or equation 46 of Sheth 1995). Iterating this process shows that

$$\begin{aligned} P(b, S_m = k) &= \sum_{l_1=1}^{k-m+1} \eta(l_1, b) P(b, S_{m-1} = k-l_1) \\ &= \sum_{l_1=1}^{k-m+1} \eta(l_1, b) \times \\ & \quad \sum_{l_2=1}^{k-l_1-m+2} \eta(l_2, b) P(b, S_{m-2} = k-l_1-l_2) \end{aligned}$$

$$\begin{aligned} &= \sum_{\text{all } m \text{ parts.}} \eta(l_1, b) \eta(l_2, b) \dots \eta(l_m, b) \\ &= \sum_{\text{all } m \text{ parts.}} \eta(1, b)^{n_1} \dots \eta(k, b)^{n_k}, \end{aligned} \quad (\text{B5})$$

where $n_1 + \dots + n_k = m$, and $l_1 + \dots + l_m = \sum_{j=1}^k j n_j = k$, and the sum in the two final expressions is over all distinct ordered partitions of k that have exactly m parts. Notice that not all the terms in this sum are different. For example, when $m = 3$ and $k = 6$, then the set $\{123\}$ occurs six times, and when $m = 3$ and $k = 7$, then the set $\{223\}$ occurs thrice. In general, a given set $\{n_1, \dots, n_k\}$, will occur $m!/(n_1! \dots n_k!)$ times.

Let $p(n_1 \dots n_k | m, k)$ denote the probability that a given set $\{n_1, \dots, n_k\}$ occurred, given that there were exactly m terms which added up to k . Then $p(n_1 \dots n_k | m, k)$ is given by summing up all the terms in equation (B5) corresponding to it, and normalizing by $P(b, S_m = k)$. If it is not certain that there were exactly m terms in the partition, then we must multiply $p(n_1 \dots n_k | m, k)$ by the probability that there were exactly m terms that added up to k . Thus, $p(n_1 \dots n_k | k)$, which is conditioned on k only and not on m as well, is given by an expression like

$$\frac{\eta(1, b)^{n_1} \eta(2, b)^{n_2} \dots \eta(k, b)^{n_k}}{P(b, S_m = k)} \frac{m!}{n_1! \dots n_k!} \times n(m|k),$$

where $n(m|k)$ denotes the probability that k is the sum of exactly m integers. Now, Appendix A showed that the branching process we are considering requires $n(m|k)$ to have the Binomial distribution of equation (7). Thus, the probability that a given set $\{n_1, \dots, n_k\}$ occurs is

$$\begin{aligned} p(n_1 \dots n_k | k) &= \frac{\eta(1, b_1)^{n_1} \dots \eta(k, b_1)^{n_k}}{P(b_1, S_m = k)} \frac{m!}{n_1! \dots n_k!} \\ & \quad \times \binom{k-1}{m-1} \left(\frac{b_1}{b_2}\right)^{k-m} \left(1 - \frac{b_1}{b_2}\right)^{m-1} \\ &= \frac{k!}{m!} \frac{(b_2 - b_1)^{m-1}}{b_2^{k-1}} \frac{e^{kb_1}}{k^{k-m}} \\ & \quad \times \frac{m!}{n_1! \dots n_k!} \prod_{j=1}^k \eta(j, b_1)^{n_j}. \end{aligned} \quad (\text{B6})$$

Simple algebra shows that this final expression is equivalent to equation (5). This shows explicitly how the partition function is related to sums of Borel-distributed random variables.

We are finally in a position to show that equation (4) follows from equation (5). The probability that a particle which is chosen at random from a clump with exactly k particles at the epoch b_2 was in a subclump with j particles at the epoch b_1 is

$$\sum_{\text{all part.}} \frac{j n_j}{k} p(n_1, \dots, n_k | k),$$

where the sum is over all partitions of k . This sum can be written as

$$\begin{aligned} & \sum_{\text{all part.}} \frac{j}{k} \eta(j, b_1) \frac{\eta(1, b_1)^{n_1} \dots \eta(j, b_1)^{n_{j-1}} \dots \eta(k, b_1)^{n_k}}{P(b_1, S_m = k)} \\ & \quad \times \frac{(m-1)!}{n_1! \dots (n_j-1)! \dots n_k!} m n(m|k), \end{aligned} \quad (\text{B7})$$

which is the same as

$$\sum_{m=1}^{k-j+1} \frac{j}{k} \eta(j, b_1) \frac{P(b_1, S_{m-1} = k-j)}{P(b_1, S_m = k)} m n(m, b_1 | k, b_2), \quad (\text{B8})$$

where it is understood that when $m = 1$ then $P(b_1, S_{m-1} = k-j) = P(b_1, S_0 = 0) = 1$, and $P(b_1, S_0 = k-j) = 0$ for $j \neq k$. Writing all the terms in the sum explicitly gives

$$\begin{aligned} & \sum_{m=1}^{k-j+1} \frac{j}{k} \frac{(jb_1)^{j-1} e^{-jb_1}}{j!} \frac{m-1}{k-j} \frac{[(k-j)b_1]^{k-j-m+1} e^{-(k-j)b_1}}{(k-j-m+1)!} \\ & \times \frac{k}{m} \frac{(k-m)!}{(kb_1)^{k-m} e^{-kb_1}} m \binom{k-1}{m-1} \left(\frac{b_1}{b_2}\right)^{k-m} \left(1 - \frac{b_1}{b_2}\right)^{m-1} \\ & = \sum_{m=2}^{k-j+1} \binom{k}{j} \frac{j^j}{k^k} \left(\frac{b_1}{b_2}\right)^{j-1} k \left(1 - \frac{b_1}{b_2}\right) \times \\ & \quad \binom{k-j-1}{m-2} \left[k \left(1 - \frac{b_1}{b_2}\right)\right]^{m-2} \left[(k-j) \frac{b_1}{b_2}\right]^{k-j-m+1}. \quad (\text{B9}) \end{aligned}$$

The Binomial theorem reduces this final expression to equation (4). This shows explicitly that

$$f(j, b_1 | k, b_2) = \sum_{\text{all part.}} \frac{j n_j}{k} p(n_1, \dots, n_k | k) \quad (\text{B10})$$

as expected. Thus, the merger probabilities of equation (4) can be derived directly from the partition function (equation 5), which means that the merger probabilities and the partition function are mutually consistent.

The steps leading to equation (B10) imply that the mean number of subclumps each having exactly j particles that are incorporated in a clump with exactly k particles is

$$\sum_{\text{all part.}} n_j p(n_1, \dots, n_k | k) = \frac{k}{j} f(j, b_1 | k, b_2). \quad (\text{B11})$$

This is the same result as that obtained using a different argument (equation 45 in Sheth 1995). However, with the partition function, it is now possible to compute the higher order moments of this distribution as well. For completeness the factorial moments are computed below. These higher order moments are useful for estimating the scatter around the mean number of j -clumps per k -clump. Also, they may be good discriminators between different partition functions that yield the same $f(j|k)$ merger probabilities. The factorial moments are

$$\begin{aligned} \left\langle \frac{n_j!}{(n_j - i)!} \right\rangle &= \sum_{m=i+1}^{k-ij+i} \eta(j, b_1)^i \frac{P(b_1, S_{m-i} = k-ij)}{P(b_1, S_m = k)} \\ &\quad \times \frac{m!}{(m-i)!} n(m, b_1 | k, b_2) \\ &= \frac{k!}{j^i (k-ij)!} \frac{(j^{j-1})^i}{k^{k-1}} \left(\frac{b_1}{b_2}\right)^{ij-i} \left[k \left(1 - \frac{b_1}{b_2}\right)\right]^i \\ &\quad \times \left[k \left(1 - \frac{b_1}{b_2}\right) + (k-ij) \frac{b_1}{b_2}\right]^{k-ij-1}. \quad (\text{B12}) \end{aligned}$$

Since the partition function is known, it is also straightforward to compute ‘cross-correlation’ type moments of the form $\langle n_i n_j \rangle$, and the associated factorial moments, though we have not done so here.

APPENDIX C: MERGING, BRANCHING, AND THE THEORY OF QUEUES

The Borel distribution (eq. 1) also arises in studies of the distribution of waiting times in queues (Borel 1942; Tanner 1953; Tanner 1961). The fact that queueing theories and the Galton–Watson branching process are closely related (Kendall 1951) will be exploited in this Appendix.

Consider a counter at which customers are served. Assume that customers arrive at the counter in a Poisson process with parameter unity (an average of one arrival per unit time) and that the service time is the same constant, $0 \leq b \leq 1$, for each customer. If the counter is busy when a customer arrives, the customer joins the back of a queue. So, service is on a first-come-first-served basis. In such a system, we can ask for the probability that exactly N customers are served before the queue is first emptied, given that there was only one customer in the queue initially. This probability is given by the Borel distribution with parameter b (Borel 1942; Tanner 1953; Consul 1989). The relation of this queue to the Galton–Watson branching process described above is clear. Simply view the N customers that were served before the queue was first emptied as the descendants, the ones who were ‘infected’ by, the initial customer.

Now modify the queue system as follows. Assume, as before, that customers arrive at a counter in a Poisson process with unit rate, and that the service time is the same constant, say b_2 , for each customer. However, in this case, the customers form two queues in accordance with the following prescription. If a new customer arrives within a time $b_1 \leq b_2$ of the most recent commencement of service, they join the back of a high priority queue, H . If they arrive after later than b_1 of the most recent commencement of service, they join the back of a low priority queue, L . All customers in queue H are serviced, in the order in which they arrived, until the queue is empty. When this happens, the first customer in queue L moves into queue H and is serviced immediately. Thus, while customers within each queue are serviced on a first-come-first-served basis, it is possible for some customers in queue H to receive service before others in queue L , even though they may have arrived later. In this sense, the modified system does not operate on a strictly first-come-first-served basis.

For such a system, we can ask for the probability that exactly k customers are served before both queues are completely emptied for the first time, given that there was only one customer in queue H and none in queue L initially. Clearly, the answer to this question is no different than before: this probability must be given by the Borel distribution with parameter b_2 . However, in the time before both queues were emptied for the first time, having serviced exactly k customers, queue H may have been emptied a number of times (though certainly not more than k times). Suppose that it was emptied m times. Define a batch of customers as the number, l , of customers served between two successive empty periods of queue H . Then we can ask for the probability, $p(l_1, l_2, \dots, l_m | k)$, that queue H was emptied m times, and that customers were served in batches of $l_1, l_2, \dots, l_m$ (not necessarily in that order), given that $l_1 + l_2 + \dots + l_m = k$, and that there was only one customer in queue H initially. Then $p(l_1, l_2, \dots, l_m | k)$ is the same as that defined in the Galton–Watson process considered ear-

lier in this paper. So, for this queue system, it is given by equation (5).

In the context of this queue system, equation (5), $\eta(l, b)$ is the Borel distribution with parameter b (eq. 1), and n_j denotes the number of times exactly j customers passed through queue H before it was emptied, given that k passed through it before both H and L were emptied. The first term on the right of equation (5) accounts for the fact that the probability of serving exactly l_i customers between two successive empty periods in the H queue is a Borel distribution, and so it weights the occurrence of each l_i in the sequence of service batches with the probability, $\eta(l_i, b_1)$, that l_i occurred. The second term on the right accounts for the different permutations of the different service sequences, since the various permutations of $\{l_1, \dots, l_m\}$ all contribute to the same $p(l_1, l_2, \dots, l_m | k)$. The final term looks like a Poisson weighting term, and is obtained by an argument that is similar to that used by Tanner (1961) in his elegant derivation of the Borel distribution.

Consider the case in which both queues are empty, having serviced k customers since the last time they were both empty. If the H queue was emptied m times during this busy period, then $m - 1$ customers must have passed through queue L during this time. So, we can ask for the probability that $m - 1$ customers arrived at and passed through queue L during this time. However, in Tanner's (1961) language, not all possible arrival patterns are 'admissible', since the arrivals in queue L must be consistent with the pattern of service times in queue H . Namely, during each of the first $m - 1$ instants at which queue H is emptied, there must be at least one customer in queue L , but the m th time that H is emptied, queue L must also be empty. The problem is to list all possible sequences of arrivals at queue L , and then to determine the fraction of these that are admissible.

Recall that customers arrive in a Poisson process with unit rate at a queueing system that has constant service time $b_2 \leq 1$ per customer, and the first customer joins queue H without passing through queue L . If a customer arrives during the first fraction, b_1/b_2 , of the service time, then they join queue H , otherwise they join queue L . If k customers were served in total, then there were k opportunities for customers to join queue L . Each of these k opportunities was of duration $b_2 - b_1$. Since the arrival of customers is random, the probability that $m - 1$ customers passed through queue L during the time in which k customers were served is given by the Poisson distribution, $[k(b_2 - b_1)]^{m-1} e^{-k(b_2 - b_1)} / (m - 1)!$. However, the probability that the $m - 1$ arrivals were in an admissible order introduces an additional factor of $1/m$. This sets the final term in equation (5).

In terms of the Galton–Watson branching process considered in the main text (in which the probability that any parent has n_1 daughters is a Poisson distribution with parameter b_1 , and the probability that that parent also had n_2 sons is a Poisson distribution with parameter $b_2 - b_1$), equation (5) lists the probability that a family of size k is made of the m subfamilies (each with a male at the head and only females in the subsequent generations), having $l_1, l_2, \dots, l_m$ members in each. So, for the gravitational clustering process, the probability that a clump of size k at the epoch b_2 had the m progenitors $l_1, l_2, \dots, l_m$ at the epoch b_1 is given by equation (5). As noted in section 3, equation (5) can be used to generate the partition function of merger history trees.

This way of deriving the partition function, by generalizing Tanner's (1961) argument, has another connection to previous work. In effect, it is an alternative derivation of the excursion set scaling solution derived in section 3.2 of Sheth (1995). This follows because Tanner showed how his queueing system could be formulated in terms of an excursion set process. Here we have described a queueing system that is associated with the Poisson Galton–Watson branching process. Comparison of this queue with the excursion set process considered in section 3.2 of Sheth (1995) shows that they are equivalent.

APPENDIX D: COUNTS-IN-CELLS FROM THE GALTON–WATSON BRANCHING PROCESS

The branching process extension of the Press–Schechter approach is very powerful. In the main text it was used to provide a description of the merger history tree. However, following a calculation suggested by Consul (1989), it is also relatively straightforward to use the Galton–Watson branching process to provide a derivation of the distribution of counts in randomly placed cells, at any epoch characterized by b . Assume that Press–Schechter Borel clumps collapse completely to points, so that if the cluster center is included in a cell, all associated particles are also. Now assume that all clumps evolve in accordance with the Press–Schechter description developed in the main text.

The analogy with the Galton–Watson process is as follows. The number of particles in a given cell is the same as the total number of progeny of the Galton–Watson process. However, unlike the Press–Schechter mass function considered in the main text, in this case the number of initial ancestors, X_0 , is not unity. Rather, it is the number of cluster centers, $X_0 = m$, say, that happen to be in the cell. So, for a cell containing m cluster centers, rather than considering the total number of progeny of one ancestor (the Borel distribution with parameter b), we need to calculate the total number of progeny given that there are $X_0 = m$ ancestors in the zeroth generation. This is the same as the Poisson Galton–Watson process, conditioned on the number of ancestors being $X_0 = m$, rather than unity. Both the branching process (Consul 1989) and the queueing theory (Tanner 1953) formulations of this problem show that the probability that there are exactly N particles in the cell given that there are exactly m cluster centers in the cell is

$$P(N|m) = \frac{m}{N} \frac{(Nb)^{N-m} e^{-Nb}}{(N-m)!}, \quad (\text{D1})$$

which is the Borel–Tanner distribution of Appendix C. When $m = 1$ it reduces to the Borel distribution of equation (1).

The final key idea is to assume that, since the initial distribution is Poisson, any randomly placed cell will include a random number X_0 of cluster centers. That is, the distribution of X_0 will be Poisson, with a parameter that is specified by the epoch, labelled by b , at which the cell is placed, and by the size of the cell. This model for the number of cluster centers in a randomly placed cell, given that the initial distribution was Poisson, is consistent with linear theory (e.g. Peebles 1980), and is motivated by a scaling argument that can be applied to the Poisson Press–Schechter description

(section 3.2 in Sheth 1995). However, the scaling argument is equivalent to the queueing theory interpretation of the Galton–Watson partition function of merger history trees (see discussion at the end of Appendix C). Thus, the assumption that the number of clusters in a randomly placed cell is a Poisson random variable is consistent with the partition structure derived earlier in this paper.

When the distribution of X_0 is Poisson with parameter $\bar{N}_{\text{cl}}$, then the probability that any randomly placed cell contains exactly N particles is

$$\begin{aligned} f(N) &= \sum_{m=0}^{\infty} p(m) P(N|m) = \sum_{m=0}^{\infty} \frac{\bar{N}_{\text{cl}}^m e^{-\bar{N}_{\text{cl}}}}{m!} P(N|m) \\ &= \frac{\bar{N}_{\text{cl}}}{N!} \sum_{m=1}^{\infty} \binom{N-1}{m-1} \bar{N}_{\text{cl}}^{m-1} (Nb)^{N-m} e^{-\bar{N}_{\text{cl}} - Nb} \\ &= \frac{\bar{N}_{\text{cl}}}{N!} (\bar{N}_{\text{cl}} + Nb)^{N-1} e^{-\bar{N}_{\text{cl}} - Nb}. \end{aligned} \quad (\text{D2})$$

Setting $\bar{N}_{\text{cl}} = \bar{n}V(1-b)$, where $\bar{n}$ is the average density of particles and V is the size of the cell, is required by normalization. This shows that equation (D2) implies that the probability that a randomly placed cell of size V contains exactly N particles is

$$f(N, V) = \frac{\bar{N}(1-b)}{N!} \left[\bar{N}(1-b) + Nb \right]^{N-1} e^{-\bar{N}(1-b) - Nb}, \quad (\text{D3})$$

where the left hand side now shows the volume dependence explicitly.

Equation (D3) is also a solution to the Saslaw & Hamilton (1984) thermodynamic model of nonlinear gravitational clustering. It can be understood as describing a Poisson distribution of point sized clumps, where the probability a clump has N associated particles is given by a Borel distribution with parameter b (Saslaw 1989). Thus, equations (1) and (D3) can be derived from a thermodynamic model (Saslaw & Hamilton 1984; Sheth 1995), from an analysis of the excursion set statistics of overdense regions of a Poisson distribution (Sheth 1995), and from the Poisson Galton–Watson branching process (Consul 1989).

In equation (D3), b is constant, independent of cell size V . This is a consequence of assuming that all clumps are point sized. Relaxing this assumption means that the branching process is no longer straightforward to implement. Nevertheless, we can use the Poisson cluster interpretation of this branching process derivation of $f(N, V)$ to gain some understanding of the shape of the counts in cells distribution when the point sized approximation is relaxed. The counts in cells distribution of a Poisson distribution of Borel (with parameter b) Press–Schechter clumps that have nontrivial sizes and shapes is well approximated by equation (D3), except that b becomes scale dependent. It tends to zero as $V \rightarrow 0$ and it tends to a constant value as V becomes larger than the typical clump size (Sheth & Saslaw 1994).

Whereas the usual Press–Schechter analysis of excursion set mass functions provides information about the distribution of virialized clump masses, it does not provide information about the internal structure of these clumps, nor does it describe how these clumps are distributed relative to each other in space. The clumps may be correlated with each other, or distributed uniformly at random. Thus, one cannot compute the N -point correlation functions of

the clustered distribution, nor can one construct the nonlinear counts in cells distribution function. In this respect, the Press–Schechter description does not provide a complete description of nonlinear clustering. Therefore, it is very interesting that numerical simulations of gravitational clustering from an initially Poisson distribution confirm the accuracy of equation (D3) on all scales, as well as the scale dependence and temporal evolution of b (Sheth & Saslaw 1994 and references therein). This measured accuracy of equation (D3), and the way in which it can be derived from the branching process, suggests one way in which the Press–Schechter approach may be extended to provide some information about the counts in cells distribution.

Before concluding, we note that the partition structure derived in the main text is independent of the accuracy or applicability of the results of this Appendix. That is, the results of the main text are independent of whether or not the Borel clusters have a Poisson spatial distribution.